

ANAPHYLAXIS EMERGENCY GAME PLAN

_____ is committed to the health, safety and wellbeing of our partners, clients and visitors. Our Code Blue Team has been trained to provide emergency first aid care in the event someone suffers an anaphylactic reaction.

Location of EpiPen: _____ has placed EpiPen & EpiPen Jr. strategically throughout the facility for emergency use in the following locations

EpiPen (For patients weighing ≥ 30 kg/66lbs or ≥ 8 years old)	EpiPen Jr (For patients weighing < 30 kg/66lbs or < 8 years old)
Insert Locations here	Insert Locations here

WHEN TO ADMINISTER EpiPen...

Confirmed or suspected exposure to allergen One (1) Airway symptom OR Two (2) Non-Airway symptoms	
Airway	Non-Airway
<ul style="list-style-type: none"> • Shortness of breath • Difficulty breathing • Coughing • Hoarse voice • Drooling • Swelling to face (tongue or lips) 	<ul style="list-style-type: none"> • Rash • Hives • Itchiness • Vomiting • Diarrhea • Abdominal Cramps
Anaphylaxis can occur up to 6 hours after exposure When in doubt give EpiPen	

Emergency Action Plan

1. Activate Emergency First Aid Game Plan. Call 9-1-1. Bring AED and Emergency Response Kit to the person including EpiPen
2. Administer the EpiPen
 - a. Pull off the Grey cap
 - b. Keeping your finger away from the blue section, inject the EpiPen into the outer thigh (Clothing should be removed)
 - c. Hold the EpiPen into the thigh (bare thigh whenever possible) for 10 – 15 seconds
3. Monitor and document the situation. If no improvement in 5 minutes administer a second EpiPen in the other thigh
4. Transfer care to paramedics upon their arrival

APPROVED DATE: _____ PLAN EXPIRY DATE: _____